

Current Status of Biological Weed Control Agents in Idaho, Oregon, and Washington

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
Bindweed, field (<i>Convolvulus arvensis</i>)	<i>Aceria malherbae</i>	L	M	F	L	L	M	F	L	L	L	F	L
	<i>Tyta luctuosa</i>	L	L	P	U	U	U	U	U	L	L	U	L
Broom, French (<i>Genista monspessulana</i>)	<i>Bruchidius villosus</i>	W	H	G	M	—	—	—	—	—	—	—	—
Broom, Scotch (<i>Cytisus scoparius</i>)	<i>Bruchidius villosus</i>	W	H	G	M	W	H	G	M	L	U	U	U
	<i>Exapion fuscirostre</i>	W	H	G	M	W	M	G	L	L	U	U	U
	<i>Leucoptera spartifoliella</i>	W	M	P	L	L	M	P	L	—	—	—	—
Crupina, common (<i>Crupina vulgaris</i>)	<i>Ramularia crupinae*</i>	U	U	U	U	—	—	—	—	U	U	U	U
Gorse (<i>Ulex europaeus</i>)	<i>Exapion ulicis</i>	W	H	G	M	W	M	F	L	—	—	—	—
	<i>Sericothrips staphylinus*</i>	L	U	U	L	L	U	U	U	—	—	—	—
	<i>Tetranychus lintearius</i>	W	L	P	L	W	L	P	L	—	—	—	—
Hawkweeds (mouse-ear [<i>Pilosella officinarum</i>]; orange [<i>P. aurantiaca</i>]; whiplash [<i>P. flagellaris</i>])	<i>Aulacidea subterminalis*</i>	—	—	—	—	U	U	U	U	—	—	—	—
	<i>Cheilosia urbana*</i>	—	—	—	—	—	—	—	—	—	—	—	—
Hemlock, poison (<i>Conium maculatum</i>)	<i>Agonopterix alstroemeriana</i>	W	L	P	M	W	L	P	L	L	L	P	L
Knapweed, diffuse (<i>Centaurea diffusa</i>)	<i>Agapeta zoegana</i>	W	U	U	L	L	U	U	L	U	U	U	U
	<i>Bangastermus fausti</i>	W	H	G	M	W	H	G	M	L	L	F	L
	<i>Chaetorellia acrolophi</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Cyphocleonus achates</i>	W	L	F	M	W	M	G	L	L	M	G	M
	<i>Larinus minutus</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Larinus obtusus</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Metzneria paucipunctella</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Pterolonche inspersa</i>	L	L	L	U	F	—	—	—	F	—	—	—
	<i>Sphenoptera jugoslavica</i>	W	H	G	L	W	H	G	M	W	H	G	M
	<i>Terellia virens</i>	U	U	U	U	U	U	U	U	U	U	U	U
continued Knapweed, diffuse (<i>Centaurea diffusa</i>)	<i>Urophora affinis</i>	W	H	G	M	W	H	G	M	W	H	G	M
	<i>Urophora quadrifasciata</i>	W	H	G	M	W	H	G	M	W	H	G	M
Knapweed, meadow	<i>Larinus minutus</i>	L	M	F	L	L	M	F	L	—	—	—	—

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
<i>(Centaurea x moncktonii; = C. debeauxii subsp. grandiflora; C. jacea nothosubsp. Pratensis)</i>	<i>Larinus obtusus</i>	W	H	G	M	W	H	G	M	—	—	—	—
	<i>Urophora quadrifasciata</i>	W	L	F	L	L	M	F	L	—	—	—	—
Knapweed, Russian <i>(Rhaponticum repens; = Acroptilon repens)</i>	<i>Aulacidea acroptilonica</i>	W	H	F	M	W	H	G	M	W	M	F	M
	<i>Jaapiella ivannikovi</i>	W	L	P	M	W	L	F	M	L	M	F	L
	<i>Mesoanguina picridis</i> (formerly <i>Subanguina picridis</i>)	L	U	U	U	F	—	—	—	U	U	U	U
Knapweed, squarrose <i>(Centaurea virgata subsp. squarrosa; = C. virgata; = C. squarrosa)</i>	<i>Bangasterus fausti</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Cyphocleonus achates</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Larinus minutus</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Sphenoptera jugoslavica</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Urophora affinis</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Urophora quadrifasciata</i>	—	—	—	—	—	—	—	—	U	U	U	U
Knapweed, spotted <i>(Centaurea stoebe; = C. stoebe ssp. micranthos)</i>	<i>Agapeta zoegana</i>	L	H	G	U	L	M	U	L	W	M	U	L
	<i>Bangasterus fausti</i>	W	L	G	M	L	L	U	L	L	U	U	L
	<i>Chaetorellia acrolophi</i>	L	L	F	L	U	U	U	U	L	L	U	L
	<i>Cyphocleonus achates</i>	W	M	F	M	L	M	F	L	W	M	G	M
	<i>Larinus minutus</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Larinus obtusus</i>	W	H	E	M	W	H	E	M	L	M	F	L
	<i>Metzneria paucipunctella</i>	W	L	F	L	W	L	F	L	W	H	G	M
	<i>Sphenoptera jugoslavica</i>	W	H	G	L	W	H	G	M	L	L	P	L
	<i>Terellia virens</i>	L	M	G	L	U	U	U	U	L	U	U	U
	<i>Urophora affinis</i>	W	M	G	M	W	H	G	M	W	H	G	M
<i>Urophora quadrifasciata</i>	W	H	G	M	W	H	G	M	W	H	G	M	
Knotweed, giant <i>(Reynoutria sachalinensis; = Polygonum sachalinense; = Fallopia sachalinensis)</i>	<i>Aphalara itadori*</i> – Hokkaido strain	U	U	U	L	U	U	U	L	—	—	—	—
Knotweed, Japanese <i>(Reynoutria japonica; = Polygonum cuspidatum; = Fallopia japonica);</i> Bohemian <i>(Reynoutria x bohemia; = Polygonum x bohemicum; = Fallopia x bohemia)</i>	<i>Aphalara itadori*</i> – Kyushu strain	L	U	U	L	U	U	U	L	U	U	U	L
Knotweed, Japanese <i>(Reynoutria japonica; = Polygonum cuspidatum; = Fallopia japonica);</i> Bohemian <i>(Reynoutria x bohemia; =</i>	<i>Aphalara itadori*</i> – Murakami strain	U	U	U	L	U	U	U	L	—	—	—	—

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
<i>Polygonum x bohemicum</i> ; = <i>Fallopia x bohémica</i>)													
Loosestrife, purple (<i>Lythrum salicaria</i>)	<i>Galerucella californiensis</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Galerucella pusilla</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Hylobius transversovittatus</i>	W	H	G	L	L	M	F	L	L	L	F	L
	<i>Nanophyes marmoratus</i>	W	M	F	M	W	M	U	L	W	L	F	M
Puncturevine (<i>Tribulus terrestris</i>)	<i>Microlarinus lareynii</i> *	L	L	P	L	L	S	P	U	L	L	P	U
	<i>Microlarinus lypriformis</i> *	F	—	—	—	U	U	U	U	L	L	P	U
Ragwort, tansy (<i>Jacobaea vulgaris</i> ; = <i>Senecio jacobaea</i>)	<i>Botanophila seneciella</i>	W	S	P	L	W	M	F	M	L	L	F	U
	<i>Longitarsus jacobaeae</i>	W	H	E	M	W	H	E	M	W	H	G	L
	<i>Tyria jacobaeae</i> *	W	H	E	N	W	H	G	N	—	—	—	N
Rush, flowering (<i>Butomus umbellatus</i>)	<i>Bagous nodulosus</i> *	—	—	—	—	—	—	—	—	—	—	—	—
Sage, Mediterranean (<i>Salvia aethiopsis</i>)	<i>Phrydiuchus tau</i>	W	H	G	L	L	M	L	N	W	H	G	M
St. Johnswort (<i>Hypericum perforatum</i>)	<i>Agrilus hyperici</i>	L	H	E	L	L	M	U	L	W	H	E	M
	<i>Aplocera plagiata</i>	W	M	P	M	L	M	U	L	W	M	F	M
	<i>Chrysolina hyperici</i>	W	H	G	L	W	H	E	L	W	H	E	M
	<i>Chrysolina quadrigemina</i>	W	H	G	L	W	H	E	L	W	H	E	M
Saltcedar (<i>Tamarix chinensis</i> , <i>T. parviflora</i> , <i>T. ramosissima</i>)	<i>Diorhabda</i> spp.*	W	M	F	L	U	U	U	U	L	H	G	L
Skeletonweed, rush (<i>Chondrilla juncea</i>)	<i>Aceria chondrillae</i> (formerly <i>Eriophyes chondrillae</i>)	W	H	E	M	W	H	G	M	W	H	E	M
	<i>Bradyrrhoa gilveolella</i>	L	S	U	U	U	U	U	U	L	M	G	L
	<i>Cystiphora schmidtii</i>	W	H	G	M	W	H	G	M	W	H	E	M
	<i>Puccinia chondrillina</i> *	W	M	G	U	W	H	G	U	W	H	G	U
Spurge, leafy (<i>Euphorbia virgata</i> ; = <i>E. esula</i>)	<i>Aphthona cyparissiae</i>	L	H	E	L	L	M	F	L	L	L	G	M
	<i>Aphthona czwalinai</i>	L	H	E	M	L	L	U	U	L	L	G	M
	<i>Aphthona flava</i>	L	H	F	L	L	M	F	L	L	L	G	M
	<i>Aphthona lacertosa</i>	W	H	E	M	L	M	U	L	W	H	E	M
	<i>Aphthona nigricutis</i>	W	H	E	M	L	M	F	U	W	H	E	M
	<i>Hyles euphorbiae</i>	F	—	—	—	—	—	—	—	W	M	G	M
	<i>Oberea erythrocephala</i>	W	L	G	M	L	U	U	U	W	H	E	M

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
Starthistle, yellow (<i>Centaurea solstitialis</i>)	<i>Bangasternus orientalis</i>	W	L	P	L	W	L	P	L	W	L	P	M
	<i>Ceratopion basicorne*</i>	—	—	—	—	—	—	—	—	U	U	U	L
	<i>Chaetorellia australis</i>	W	H	E	M	W	M	G	M	W	M	G	M
	<i>Eustenopus villosus</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Larinus curtus</i>	W	H	E	N	W	M	G	LN	W	H	E	N
	<i>Puccinia jacea</i> var. <i>solstitialis*</i>	L	S	U	U	U	U	U	U	U	U	U	U
	<i>Urophora sirunaseva</i>	W	L	P	L	W	L	U	L	W	L	U	L
Thistle, bull (<i>Cirsium vulgare</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	U	U	L	W	M	L	U	U	U	U	L
	<i>Rhinocyllus conicus*</i>	W	M	F	N	W	L	L	N	W	M	F	N
	<i>Trichosirocalus horridus*</i>	L	U	U	N	U	U	U	N	L	L	U	N
	<i>Urophora stylata</i>	W	H	G	M	L	M	F	L	L	L	U	L
Thistle, Canada (<i>Cirsium arvense</i>)	<i>Hadroplontus litura</i> (formerly <i>Ceutorhynchus litura</i>)	L	H	P	U	L	M	P	L	L	L	F	L
	<i>Puccinia punctiformis*</i>	L	H	G	U	U	U	U	U	L	L	L	U
	<i>Rhinocyllus conicus*</i>	W	H	F	N	W	H	P	N	W	F	F	N
	<i>Urophora cardui</i>	W	H	P	M	W	H	P	M	L	M	F	M
Thistle, Italian (<i>Carduus pycnocephalus</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	H	L	L	—	—	—	N	—	—	—	N
Continued Thistle, Italian (<i>Carduus pycnocephalus</i>)	<i>Rhinocyllus conicus*</i>	W	H	G	N	—	—	—	N	L	H	G	N
	<i>Trichosirocalus horridus*</i>	W	H	U	N	—	—	—	N	L	L	U	N
Thistle, milk (<i>Silybum marianum</i>)	<i>Rhinocyllus conicus*</i>	W	H	G	N	—	—	—	N	—	—	—	—
Thistle, musk (<i>Carduus nutans</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	U	U	L	U	U	U	U	U	U	U	U
	<i>Rhinocyllus conicus*</i>	W	H	G	N	W	H	G	N	W	H	G	N
	<i>Trichosirocalus horridus*</i>	L	M	G	N	—	—	—	N	L	M	G	N
Thistle, plumeless (<i>Carduus acanthoides</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	—	U	U	U	U	U	U	U	U	U	U	U
	<i>Rhinocyllus conicus*</i>	L	H	G	N	W	H	G	N	W	H	G	N
	<i>Trichosirocalus horridus*</i>	U	U	U	N	L	L	U	N	L	L	U	N
Thistle, slenderflower (<i>Carduus tenuiflorus</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	M	F	M	L	U	U	N	—	—	—	—
	<i>Rhinocyllus conicus*</i>	W	H	G	N	—	—	—	N	—	—	—	—

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
	<i>Trichostirocalus horridus</i> *	W	H	F	N	—	—	—	N	—	—	—	—
Toadflax, Dalmatian (<i>Linaria dalmatica</i>)	<i>Brachypterosus pulicarius</i>	W	M	P	M	W	M	P	M	W	H	E	M
	<i>Calophasia lunula</i>	L	L	P	U	W	M	P	U	W	L	U	L
	<i>Mecinus janthiniformis</i> (formerly <i>M. janthinus</i>)	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Rhinusa</i> spp.	W	U	U	L	W	U	U	L	U	U	U	U
Toadflax, yellow (<i>Linaria vulgaris</i>)	<i>Brachypterosus pulicarius</i>	L	M	F	L	L	M	F	L	L	M	P	L
	<i>Calophasia lunula</i>	L	U	U	U	L	U	U	U	L	M	F	L
	<i>Rhinusa</i> spp.	W	M	U	L	L	M	U	L	L	M	U	L
	<i>Rhinusa pilosa</i> *	U	U	U	U	—	—	—	—	U	U	U	L
	<i>Mecinus janthinus</i>	L	H	E	L	L	M	U	U	W	H	E	L
Watermilfoil, Eurasian (<i>Myriophyllum spicatum</i>)	<i>Euhrychiopsis lecontei</i> *	L	L	U	U	W	M	F	U	W	L	F	L
Whitetop or hoary cress (<i>Lepidium draba</i>)	<i>Aceria drabae</i> *	—	—	—	—	—	—	—	—	U	U	U	L

¹ Distribution within host range: W = widespread; L = limited sites; F = failed to establish; U = unknown status; — = not released

² Attack rate host: H = heavy (> 70%); M = medium (> 30%); L = light (> 10%); S = slight (< 1%); U = unknown status

³ Control ability on seeds and/or plant density: E = excellent; G = good; F = fair; P = poor; U = undetermined

⁴ Availability for redistribution: M = mass collections; L = limited (Limited availability indicates biocontrol agent populations are slow in building or are recently introduced. Work on these species should be coordinated through biological control specialists at the state department of agriculture or state university); U = unavailable; N = not recommended. These biocontrol agents may attack native species and interstate shipments are no longer permitted.

* Indicates biocontrol agents where redistribution limitations may apply. See Table 2 for additional information.