

Current Status of Biological Weed Control Agents in Idaho, Oregon, and Washington

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
Bindweed, field (<i>Convolvulus arvensis</i>)	<i>Aceria malherbae</i>	L	M	F	L	L	M	F	M	L	L	F	L
	<i>Tyta luctuosa</i>	L	L	P	U	U	U	U	U	L	L	U	L
Broom, French (<i>Genista monspessulana</i>)	<i>Bruchidius villosus</i>	W	H	G	M	—	—	—	—	—	—	—	—
Broom, Scotch (<i>Cytisus scoparius</i>)	<i>Bruchidius villosus</i>	W	H	G	M	W	H	G	M	L	U	U	U
	<i>Exapion fuscirostre</i>	W	H	G	M	W	M	G	L	L	U	U	U
	<i>Leucoptera spartifoliella</i>	W	M	P	L	L	M	P	L	—	—	—	—
Crupina, common (<i>Crupina vulgaris</i>)	<i>Ramularia crupinae</i> *	U	U	U	U	—	—	—	—	U	U	U	U
Gorse (<i>Ulex europaeus</i>)	<i>Exapion ulicis</i>	W	H	G	M	M	H	F	L	—	—	—	—
	<i>Sericothrips staphylinus</i> *	L	U	U	L	L	U	U	L	—	—	—	—
	<i>Tetranychus lintearius</i>	W	L	P	L	W	L	P	L	—	—	—	—
Hawkweeds (mouse-ear [<i>Pilosella officinarum</i>]; orange [<i>P. aurantiaca</i>]; whiplash [<i>P. flagellaris</i>])	<i>Aulacidea subterminalis</i> *	—	—	—	—	U	U	U	U	—	—	—	—
	<i>Cheilosia urbana</i> *	—	—	—	—	—	—	—	—	—	—	—	—
Hemlock, poison (<i>Conium maculatum</i>)	<i>Agonopterix alstroemeriana</i>	W	L	P	M	W	L	P	L	L	L	P	L
Knapweed, diffuse (<i>Centaurea diffusa</i>)	<i>Agapeta zoegana</i>	W	U	U	L	L	U	U	L	U	U	U	U
	<i>Bangasternus fausti</i>	W	H	G	M	W	H	G	M	L	L	F	L
	<i>Chaetorellia acrolophi</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Cyphocleonus achates</i>	W	L	F	M	W	M	G	L	L	M	G	M
	<i>Larinus minutus</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Larinus obtusus</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Metzneria paucipunctella</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Pterolonche dispersa</i>	L	L	L	U	F	—	—	—	F	—	—	—
	<i>Sphenoptera jugoslavica</i>	W	H	G	L	W	H	G	M	W	H	G	M

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
	<i>Terellia virens</i>	U	U	U	U	U	U	U	U	U	U	U	U
	<i>Urophora affinis</i>	W	H	G	M	W	H	G	M	W	H	G	M
	<i>Urophora quadrifasciata</i>	W	H	G	M	W	H	G	M	W	H	G	M
Knapweed, meadow (<i>Centaurea x moncktonii</i> ; = <i>C. debeauxii</i> subsp. <i>grandiflora</i> ; <i>C. jacea</i> nothosubsp. <i>pratensis</i>)	<i>Larinus minutus</i>	L	M	F	L	L	M	F	L	—	—	—	—
	<i>Larinus obtusus</i>	W	H	G	M	W	H	G	M	—	—	—	—
	<i>Urophora quadrifasciata</i>	W	L	F	L	L	M	F	L	—	—	—	—
Knapweed, Russian (<i>Rhaponticum repens</i> ; = <i>Acroptilon repens</i>)	<i>Aulacidea acroptilonica</i>	W	H	F	M	W	H	G	M	W	M	F	M
	<i>Jaapiella ivannikovi</i>	W	L	P	M	W	M	F	M	L	M	F	L
	<i>Mesoanguina picridis</i> (formerly <i>Subanguina picridis</i>)	L	U	U	U	F	—	—	—	U	U	U	U
Knapweed, squarrose (<i>Centaurea virgata</i> subsp. <i>squarrosa</i> ; = <i>C. virgata</i> ; = <i>C. squarrosa</i>)	<i>Bangasternus fausti</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Cyphocleonus achates</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Larinus minutus</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Sphenoptera jugoslavica</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Urophora affinis</i>	—	—	—	—	—	—	—	—	U	U	U	U
	<i>Urophora quadrifasciata</i>	—	—	—	—	—	—	—	—	U	U	U	U
Knapweed, spotted (<i>Centaurea stoebe</i> ; = <i>C. stoebe</i> ssp. <i>micranthos</i>)	<i>Agapeta zoezana</i>	L	H	G	U	L	M	U	L	W	M	U	L
	<i>Bangasternus fausti</i>	W	L	G	M	L	L	U	L	L	U	U	L
	<i>Chaetorellia acrolophi</i>	L	L	F	L	L	L	U	L	L	L	U	L
	<i>Cyphocleonus achates</i>	W	M	F	M	L	M	F	L	L	M	G	M
	<i>Larinus minutus</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Larinus obtusus</i>	W	H	E	M	W	H	E	M	L	M	F	L
	<i>Metzneria paucipunctella</i>	W	L	F	L	W	L	F	L	W	H	G	M
	<i>Sphenoptera jugoslavica</i>	W	H	G	L	W	H	G	M	L	L	P	L
	<i>Terellia virens</i>	L	M	G	L	L	U	U	U	L	U	U	U
	<i>Urophora affinis</i>	W	M	G	M	W	H	G	M	W	H	G	M
	<i>Urophora quadrifasciata</i>	W	H	G	M	W	H	G	M	W	H	G	M
Knotweed, giant (<i>Reynoutria sachalinensis</i> ; = <i>Polygonum sachalinense</i> ; =	<i>Aphalara itadori</i> * – Hokkaido strain	U	U	U	L	U	U	U	L	—	—	—	—

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
<i>Fallopia sachalinensis</i>)													
Knotweed, Japanese <i>(Reynoutria japonica; = Polygonum cuspidatum; = Fallopia japonica);</i> Bohemian <i>(Reynoutria x bohemica; = Polygonum x bohemicum; = Fallopia x bohemica)</i>	<i>Aphalara itadori</i> * – Kyushu strain	L	U	U	L	U	U	U	L	U	U	U	L
Knotweed, Japanese <i>(Reynoutria japonica; = Polygonum cuspidatum; = Fallopia japonica);</i> Bohemian <i>(Reynoutria x bohemica; = Polygonum x bohemicum; = Fallopia x bohemica)</i>	<i>Aphalara itadori</i> * – Murakami strain	U	U	U	L	U	U	U	L	—	—	—	—
Loosestrife, purple <i>(Lythrum salicaria)</i>	<i>Galerucella californiensis</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Galerucella pusilla</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Hylobius transversovittatus</i>	W	H	G	L	L	M	F	L	L	L	F	L
	<i>Nanophyes marmoratus</i>	W	M	F	M	L	M	F	M	W	L	F	M
Puncturevine <i>(Tribulus terrestris)</i>	<i>Microtharax lareynii</i> *	L	L	P	L	L	L	P	U	L	L	P	U
	<i>Microtharax lypriformis</i> *	F	—	—	—	U	U	U	U	L	L	P	U
Ragwort, tansy <i>(Jacobaea vulgaris; = Senecio jacobaea)</i>	<i>Botanophila seneciella</i>	W	S	P	L	W	H	G	M	L	L	F	U
	<i>Longitarsus jacobaeae</i>	W	H	E	M	W	H	E	M	W	H	G	L
	<i>Tyria jacobaeae</i> *	W	H	E	M	W	H	G	M	—	—	—	N
Sage, Mediterranean <i>(Salvia aethiopis)</i>	<i>Phrydiuchus tau</i>	W	H	G	L	L	M	L	N	W	H	G	M
St. Johnswort <i>(Hypericum perforatum)</i>	<i>Agrilus hyperici</i>	L	H	E	L	L	M	U	L	W	H	E	M
	<i>Aplocera plagiata</i>	W	M	P	M	L	M	U	L	W	M	F	M
	<i>Chrysolina hyperici</i>	W	H	G	L	W	H	E	M	W	H	E	M
	<i>Chrysolina quadrigemina</i>	W	H	G	L	W	H	E	M	W	H	E	M
Saltcedar <i>(Tamarix chinensis, T. parviflora, T. ramosissima)</i>	<i>Diorhabda</i> spp.*	W	M	F	L	—	U	—	U	L	H	G	L

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
Skeletonweed, rush (<i>Chondrilla juncea</i>)	<i>Aceria chondrillae</i> (formerly <i>Eriophyes chondrillae</i>)	W	H	E	M	W	H	G	M	W	H	E	M
	<i>Bradyrrhoa gilveolella</i>	L	S	U	U	U	U	U	U	L	M	G	L
	<i>Cystiphora schmidtii</i>	W	H	G	M	W	H	G	M	W	H	E	M
	<i>Puccinia chondrillina</i> *	W	M	G	L	W	H	G	M	W	H	G	M
Spurge, leafy (<i>Euphorbia virgata</i> ; = <i>E. esula</i>)	<i>Aphthona cyparissiae</i>	L	H	E	L	L	M	F	L	L	L	G	M
	<i>Aphthona czwalinai</i>	L	H	E	M	L	L	U	U	L	L	G	M
	<i>Aphthona flava</i>	L	H	F	L	L	M	F	L	L	L	G	M
	<i>Aphthona lacertosa</i>	W	H	E	M	L	M	U	L	W	H	E	M
	<i>Aphthona nigriscutis</i>	W	H	E	M	L	M	F	U	W	H	E	M
	<i>Hyles euphorbiae</i>	F	—	—	—	—	—	—	—	W	M	G	M
	<i>Oberea erythrocephala</i>	W	L	G	M	L	U	U	U	W	H	E	M
Starthistle, yellow (<i>Centaurea solstitialis</i>)	<i>Bangasternus orientalis</i>	W	L	P	L	W	L	P	L	W	L	P	M
	<i>Ceratapion basicorne</i> *	—	—	—	—	—	—	—	—	U	U	U	L
	<i>Chaetorellia australis</i>	W	H	E	M	W	M	G	M	W	M	G	M
	<i>Eustenopus villosus</i>	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Larinus curtus</i>	W	H	E	M	W	M	F	L	W	H	E	M
	<i>Puccinia jacea</i> var. <i>solstitialis</i> *	L	S	U	U	U	U	U	U	U	U	U	U
	<i>Urophora sirunaseva</i>	W	L	P	L	W	L	U	L	W	L	U	L
Thistle, bull (<i>Cirsium vulgare</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	U	U	L	W	M	L	U	U	U	U	L
	<i>Rhinocyllus conicus</i> *	W	M	F	N	W	L	L	N	W	M	F	N
	<i>Trichosirocalus horridus</i> *	L	U	U	N	U	U	U	N	L	L	U	N
	<i>Urophora stylata</i>	W	H	G	M	L	M	F	L	L	L	U	L
Thistle, Canada (<i>Cirsium arvense</i>)	<i>Hadroplontus litura</i> (formerly <i>Ceutorhynchus litura</i>)	L	H	P	U	L	M	P	L	L	L	F	L
	<i>Puccinia punctiformis</i> *	L	H	G	U	U	U	U	U	L	L	L	U
	<i>Rhinocyllus conicus</i> *	W	H	F	N	W	H	P	N	W	F	F	N
	<i>Urophora cardui</i>	W	H	P	M	W	H	P	M	L	M	F	M

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
Thistle, Italian (<i>Carduus pycnocephalus</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	H	L	L	—	—	—	N	—	—	—	N
	<i>Rhinocyllus conicus</i> *	W	H	G	N	—	—	—	N	L	H	G	N
	<i>Trichosiocalus horridus</i> *	W	H	U	N	—	—	—	N	L	L	U	N
Thistle, milk (<i>Silybum marianum</i>)	<i>Rhinocyllus conicus</i> *	W	H	G	N	—	—	—	N	—	—	—	—
Thistle, musk (<i>Carduus nutans</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	U	U	L	U	U	U	U	U	U	U	U
	<i>Rhinocyllus conicus</i> *	W	H	G	N	W	H	G	N	W	H	G	N
	<i>Trichosiocalus horridus</i> *	L	M	G	N	—	—	—	N	L	M	G	N
Thistle, plumeless (<i>Carduus acanthoides</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	—	U	U	U	U	U	U	U	U	U	U	U
	<i>Rhinocyllus conicus</i> *	L	H	G	N	W	H	G	N	W	H	G	N
	<i>Trichosiocalus horridus</i> *	U	U	U	U	L	L	U	N	L	L	U	N
Thistle, slenderflower (<i>Carduus tenuiflorus</i>)	<i>Cheilosia grossa</i> (formerly <i>C. corydon</i>)	W	M	F	M	L	U	U	N	—	—	—	—
	<i>Rhinocyllus conicus</i> *	W	H	G	N	—	—	—	N	—	—	—	—
	<i>Trichosiocalus horridus</i> *	W	H	F	N	—	—	—	N	—	—	—	—
Toadflax, Dalmatian (<i>Linaria dalmatica</i>)	<i>Brachypterolus pulicarius</i>	W	M	P	M	W	M	P	M	W	H	E	M
	<i>Calophasia lunula</i>	L	L	P	U	W	M	P	U	W	L	U	L
	<i>Mecinus janthiniformis</i> (formerly <i>M. janthinus</i>)	W	H	E	M	W	H	E	M	W	H	E	M
	<i>Rhinusa</i> spp.	W	U	U	L	W	U	U	L	U	U	U	U
Toadflax, yellow (<i>Linaria vulgaris</i>)	<i>Brachypterolus pulicarius</i>	L	M	F	L	L	M	F	L	L	M	P	L
	<i>Calophasia lunula</i>	L	U	U	U	L	U	U	U	L	M	F	L
	<i>Rhinusa</i> spp.	W	M	U	L	L	M	U	L	L	M	U	L
	<i>Rhinusa pilosa</i> *	U	U	U	U	—	—	—	—	U	U	U	L
	<i>Mecinus janthinus</i>	L	H	E	L	L	M	U	U	W	H	E	L
Watermilfoil, Eurasian (<i>Myriophyllum spicatum</i>)	<i>Euhrychiopsis lecontei</i> *	L	L	U	U	W	M	F	U	W	L	F	L
Whitetop or hoary cress	<i>Aceria drabae</i> *	—	—	—	—	—	—	—	—	U	U	U	L

Weed	Biocontrol Agent	Oregon				Washington				Idaho			
		Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴	Distribution ¹	Attack Rate ²	Control ³	Availability ⁴
<i>(Lepidium draba)</i>													
¹ Distribution within host range: W = widespread; L = limited sites; F = failed to establish; U = unknown status; — = not released ² Attack rate host: H = heavy (> 70%); M = medium (> 30%); L = light (> 10%); S = slight (< 1%); U = unknown status ³ Control ability on seeds and/or plant density: E = excellent; G = good; F = fair; P = poor; U = undetermined ⁴ Availability for redistribution: M = mass collections; L = limited (Limited availability indicates biocontrol agent populations are slow in building or are recently introduced. Work on these species should be coordinated through biological control specialists at the state department of agriculture or state university); U = unavailable; N = not recommended. These biocontrol agents may attack native species and interstate shipments are no longer permitted. * Indicates biocontrol agents where redistribution limitations may apply. See Table 2 for additional information.													