

Crop Rotation Intervals (months) for Common Soil-active Herbicides Used in Vegetables and Other Rotational Crops*

| Herbicide | Beans (snap) | Beets (table) | Broccoli, Cauliflower & Cabbage | Carrot | Collard | Corn (sweet) | Cucumbers | Eggplant | Lettuce | Onions & Leeks | Pea | Pepper | Pumpkins & Squash | Spinach | Swiss chard | Tomato | Cereal |
|--------------|--------------------|---------------------|---------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------------------|---------------------|---------------------|---------------------|---------------------|---------------------|----------------------|
| Accent | 10 | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | 10 | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | 10 | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | 10/18 ¹ | NSP ² |
| Assure | 0 | 0 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 0 | 4 | 4 | 4 | - | 4 | - |
| Buctril | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Callisto | 18 | 18 | 18 | 18 | 18 | 0 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 3 |
| Caparol | 12 | 8 | 12 | 5 | 5 | 5 | 12 | 12 | 12 | 8 | 5 | 12 | 12 | 12 | 12 | 12 | 12 |
| Command | 0/9 | 16 | 9/16 ³ | 16 | 16 | 9/12 ⁴ | 0/9 ⁵ | 16 | 16 | 16 | 0/9 | 0/9 ⁶ | 0/9 ⁷ | 16 | 16 | 9/12 ⁸ | 12/16 ⁹ |
| Curbit | - | 8/13 ¹⁰ | - | - | - | - | 0 | - | - | - | - | - | 0 | 8/13 ¹¹ | - | - | 8 ¹² |
| Dacthal | 8 | 8 | 0 | 8 | 0 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 0 | 8 | 8 |
| Dual Magnum | 0 | 0 | NSP ¹³ | NSP | NSP | 0 | NSP | NSP | NSP | NSP | 0 | NSP | NSP | NSP | NSP | 0 | 4.5 |
| Eptam | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Goal | 2 | 2/3 ¹⁴ | 4 | 3 | 4 | 10 | 3/4 ¹⁵ | 4 | 3/4 ¹⁵ | 6 | 2 | 3/4 ¹⁵ | 3/4 ¹⁵ | 4 | 4 | 1/4 ¹⁶ | 10 |
| Impact | 9 ^{16a} | 18 | 18 | 18 | 18 | 0 | 18 | 18 | 18 | 18 | 9 | 18 | 18 | 18 | 18 | 18 | 3 |
| Karmex31 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 |
| Laudis | 10 | 18 | 18 | 18 | 18 | 0 | 18 | 18 | 18 | 18 | 10 | 18 | 18 | 18 | 18 | 10 | 4 |
| Lorox | 12 | 12 | 12 | 0 | 12 | 0 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 |
| Norton | 6/12 ¹⁷ | 0 | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 6/12 ¹⁷ | 0/12 ¹⁷ |
| Outlook | FS ¹⁸ | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | 0/4 ¹⁹ |
| Prefar | 4 ²⁰ | - | 0 | 0 | 0 | 4 ²⁰ | 0 | 0 | 0 | 4 ^{20,21} | 4 ²⁰ | 0 | 0 | 4 | 4 ²⁰ | - | 4 ²⁰ |
| Prowl | 0 | 12/14 ²² | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ | NS/24 ¹⁸ | NS/24 ²³ | NS/24 ²³ | 0 | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ | NS/24 ²³ |
| Pursuit | 4 | 40 | 40 | 40 | 40 | 18 | 40 | 40 | 18 | 40 | 4 | 40 | 40 | 40 | 40 | 40 | 4/10 ²⁴ |
| Pyramin | FS ¹⁸ | 0 | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | FS | - | FS | - |
| Quinclorac | 10 | 10 | 10 | 24 | 10 | 10 | 10 | 12 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 24 | 10 |
| Raptor | 0 | 26 | 18 | 18 | 18 | 8.5 | 18 | 18 | 9 | 9 | 0 | 18 | 9 | 18 | 18 | 18 | 3/4/18 ²⁵ |
| Reflex | 0 | 18 | 18 | 18 | 18 | 10 | 0 | 18 | 18 | 18 | 10 | 18 | 0 | 18 | 18 | 18 | 4 |
| Sandea | 9 | 24 | 18 | 15 | 18 | 3 | 9 | 12 | 18 | 18 | 9 | 10 | 9 | 24 | - | 8 | 2 |
| Sencor | 12 | 18 | 12 | 0 | 12 | 4 | 12 | 12 | 12 | 12 | 8 | 12 | 12 | 12 | 12 | 4 | 4 |
| Spartan/Zeus | 12 | 36 | 12 | 12 | 12 | 18 | 12 | 12 | 12 | 12 | 0/2 ²⁶ | 12 | 12 | 12 | 12 | 12 | 4/12 ²⁷ |
| Spin-Aid | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 |
| Stinger | 12 | 0 | 12 | 18 | 0 | 0 | 12 | 12 | 18 | 12 | 18 | 12 | 12 | 0 | 12 | 18 | 12 |
| Treflan | 0 | 12/14 ²⁸ | 0 | 0 | 0 | 5 | 5 | 5 | 5 | 5 | - | 0/5 ²⁹ | 5 | 12/14 ²⁸ | 5 | 0/5 ²⁹ | 12/14 ³⁰ |
| Upbeet | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 3 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk | 2 wk |

* Footnotes: See next page.

*** Footnotes: Crop Rotation Intervals (months) for Common Soil-active Herbicides Used in Vegetables and Other Rotational Crops**

Aim, Basagran Gramoxone, Poast, Ro-Neet, Roundup, Select, and Touchdown: No rotation restrictions listed for these crops.

¹ pH < 6.5 = 18 months

² NSP = next spring

³ cabbage 9 months if transplanted; 16 months for all direct-seeded brassica crops or >2 pints/A

⁴ corn is 9 months if <2.67 pints/A; 12 months if >2.67 pints/A

⁵ 0 months if <1.3 pints/A; 9 months if >2 pints/A

⁶ 9 months at 1.25 lb ai/A

⁷ pumpkins, 0 months if <1.3 pints/A; winter squash, 0 months if <2 pints/A; other cucurbits = 9 months

⁸ direct-seeded tomato is 12 months

⁹ wheat 12 months, all others 16 months

¹⁰ 8 months if >1.3 pints/A

¹¹ 13 months if >3 pints/A; 8 months if <3 pints/A and moldboard plowed

¹² oats

¹³ NSP = next spring

¹⁴ 3 months if >1 pint/A

¹⁵ 4 months if >1 pint/A

¹⁶ transplanted, 1 months; seeded, 2 months at 1 pint/A, 4 months >1 pint/A

^{16a} $\frac{3}{4}$ oz rate only

¹⁷ 12 months if >12 oz; 0 months for ryegrass

¹⁸ FS = following spring

¹⁹ 0 months for sorghum

²⁰ 4 months; soil must be tilled to 4 inches

²¹ 4 months on dry bulb in Willamette Valley, with soil tilled to 4 inches; 0 months in other areas

²² 12 months after spring application; 14 months after fall application

²³ NS (next spring crop) if <2 lb ai/A; otherwise, 24 months

²⁴ 4 months for wheat and rye; 10 months for barley

²⁵ 3 months for wheat; 4 months for rye; 9–18 months for barley depending on rainfall and pH

²⁶ 0 months for dry shell peas

²⁷ 4 months for wheat

²⁸ 12 months after spring application; 14 months after fall application

²⁹ 5 months before seeding; 0 months before transplanting

³⁰ 12 months after spring application; 14 months after fall application for cereals and grasses

³¹ diuron, following grass seed; 12 months following oats.