

Susceptibility of Broadleaf Weeds in Turf to Common Herbicides

Weed	2,4-D	dicamba	MCPP	MCPA	Mixtures with triclopyr	Mixtures without triclopyr
<i>Bindweed, field (Convolvulus arvensis)</i>	S-I	S	I	—	S	S
<i>Buttercup, creeping* (Ranunculus repens)</i>	S-I	S-I	I	S-I	—	I
<i>Carrot, wild (Daucus carota)</i>	I	S	I	—	—	S
<i>Chicory (Cichorium intybus)</i>	S	S	S	—	S	S
<i>Chickweed, common (Stellaria media)</i>	R	S	S-I	—	S	S
<i>Chickweed, mouse-ear (Cerastium vulgatum)</i>	I-R	S	S-I	—	S	S
<i>Clover, hop (Trifolium spp.)</i>	I	S	S-I	I	S	S
<i>Clover, subterranean (Trifolium subterraneum)</i>	I	S	I	I	S	S
<i>Clover, white (Trifolium repens)</i>	I	S	S-I	I	S	S
<i>Cranesbill (Geranium carolinianum)</i>	S-I	S	S-I	—	S	S
<i>Daisy, English** (Bellis perennis)</i>	R	I	I	—	S	S-I
<i>Daisy, oxeye (Chrysanthemum leucanthemum)</i>	I	I	I	S	—	S-I
<i>Dandelion, common (Taraxacum officinale)</i>	S	S	S-I	S-I	S	S
<i>Dandelion, false (Hypochoeris radicata)</i>	S	I	R	—	S	S
<i>Healall (Prunella vulgaris)</i>	S-I	S	S-I	—	I	S
<i>Henbit (Lamium amplexicaule)</i>	I-R	S	I	—	S	S
<i>Ivy, ground (Glechoma hederacea)</i>	I-R	S-I	I	—	S	R
<i>Knotweed, prostrate (Polygonum aviculare)</i>	I-R	S	I	—	S	S
<i>Lambsquarters (Chenopodium album)</i>	S	S	S	S	S	S
<i>Medic, black (Medicago lupulina)</i>	I-R	S-I	I	—	S	S-I
<i>Mustard, wild (Brassica kaber)</i>	S	S-I	I	S	S	S
<i>Pearlwort (Sagina procumbens)</i>	I-R	S	S	—	—	S
<i>Pigweed, prostrate (Amaranthus blitoides)</i>	S	S	S	S	S	S

<i>Pineappleweed (Matricaria matricarioides)</i>	I-R	I	I	—	—	I
<i>Plantain, broadleaf (Plantago major)</i>	S	R	I	I	S	S
<i>Plantain, buckhorn (Plantago lanceolata)</i>	S	R	I	I	S	S
<i>Purslane, common (Portulaca oleracea)</i>	I	S	R	I	S	S
<i>Shepherd's purse (Capsella bursa-pastoris)</i>	S	S	S-I	S	S	S
<i>Sorrel, red (Rumex acetosella)</i>	I-R	S	R	—	—	S
<i>Speedwells*** (Veronica spp.)</i>	R	R	R	—	R	R
<i>Spurge, prostrate (Euphorbia supina)</i>	I	S-I	I	—	S	S-I
<i>Thistle (Cirsium spp.)</i>	S-I	S	I	I	S	S
<i>Violet, wild (Viola spp.)</i>	I-R	I-R	I-R	—	S	I-R
<i>Woodsorrel, yellow (Oxalis corniculata)</i>	R	I	I-R	—	S	I-R
<i>Yarrow, common (Achillea millefolium)</i>	I	S	I-R	—	S	S-I
S = susceptible; I = intermediate (may require several successive applications for control); R = resistant *Sulfentrazone will give good control of buttercup. **Penoxsulam gives good control of English daisy. ***Mixtures with triclopyr may control <i>Veronica chamaedrys</i>. Quinclorac will control English daisies and common, slender, and thymeleaf speedwell with one application. See label for seed oil or crop oil additive rates to use with quinclorac.						